



FUNGICIDE EFFICACY FOR CORN FOLIAR DISEASES RESOURCE



Fungicide Efficacy for Control of Corn Foliar Diseases Table (02/2025)

Efficacy categories:


P = Poor; F = Fair; G = Good; VG = Very Good; E = Excellent; NL = Not labeled for use against this disease; U = Labeled, but unknown efficacy or insufficient data to rank.

Fungicide mode of action groups:

Group 11 QoI Strobilurins Group 3 DMI Triazoles Group 7 SDHI

	Active ingredient (%)	Product/Trade name	Rate/A (fl oz)	Commonly tested rate/A (fl oz)	Antracnose leaf blight	Common rust	Eyespot	Gray leaf spot	Northern corn leaf blight	Southern rust	Tar spot ¹	Harvest Restriction ²
11	Azoxystrobin 22.9%	Quadris 2.08 SC, multiple generics	6.0-15.5	6	VG	E	VG	E	G	VG	NL	7 days
	Pyraclostrobin 23.6%	Headline 2.09 EC/SC	6.0-12.0	6	VG	E	F	E	VG	VG	NL	7 days
	Picoxystrobin 22.5%	Approach 2.08 SC	3.0-12.0	6	VG	VG-E	VG	F-VG	VG	G	G ³	7 days
3	Flutriafol 20.9%	Xyway LFR 1.92 SC	LFR: 5.8-15.2	15.2	NL	U	NL	G	VG	NL	P ³	7 days
	Flutriafol 26.4%	Xyway 3D 2.5 SC	3D: 5.8-11.8	11.8								
	Propiconazole 41.8%	Tilt 3.6 EC, multiple generics	2.0-4.0	4	NL	VG	E	G	G	F	NL	30 days
	Prothioconazole 41.0%	Proline 480 SC	5.7	5.7	U	VG	E	U	VG	G	NL	14 days
	Tebuconazole 38.7%	Folicur 3.6 F, multiple generics	4.0-6.0	NA	NL	U	NL	U	VG	F	NL	36 days
	Tetraconazole 20.5%	Domark 230 ME	4.0-6.0	NA	U	U	U	E	VG	G	U	R3 (milk)
11	Azoxystrobin 13.5%	Quilt Xcel 2.2 SE, multiple generics	10.5-14.0	10.5	VG	VG-E	VG-E	E	VG	VG	U	30 days
3	Propiconazole 11.7%											
7	Benzovindiflupyr 2.9%	Triapro 2.21 SE	13.7	13.7	U	U	U	E	VG	E	G-VG	30 days
11	Azoxystrobin 10.5%											
3	Propiconazole 11.9%											
3	Cyproconazole 7.17%	Approach Prima 2.34 SC	3.4-6.8	6.8	U	U	U	E	VG	G	G-VG	30 days
11	Picoxystrobin 17.94%											
3	Flutriafol 19.3%	Fortix 3.22 SC	4.0-6.0	5	U	U	U	E	VG	VG	G-VG ³	30 days
11	Fluxasystrobin 14.84%	Preemptor 3.22 SC										
3	Flutriafol 26.74%	Lucento 4.17 SC	3.0-5.5	5	U	U	U	VG-E	VG	G-VG	G	30 days
7	Bixafen 15.55%											
3	Flutriafol 18.63%	Topguard EQ 4.29 SC	5.0-7.0	5	U	F	U	VG	G-VG	G-VG	G-VG ³	7 days
11	Azoxystrobin 25.30%											
3	Mefenfluoconazole 17.56%	Veltima 3.34 SC	7.0-10.0	7	U	U	U	VG-E	VG-E	VG-E	VG	21 days
11	Pyraclostrobin 17.56%											
3	Mefenfluoconazole 11.61%	Revytek 4.44 SC	8.0-15.0	8	U	U	U	VG-E	VG-E	VG	VG	21 days
7	Fluxapyroxad 7.74%											
11	Pyraclostrobin 15.49%											
3	Prothioconazole 16.0%	Delaro325 SC	8.0-12.0	8	VG	E	VG	E	VG	G-VG	G-VG	14 days
11	Trifloxystrobin 13.7%											
3	Prothioconazole 14.9%	Delaro Complete 3.83 SC	4.0-12.0	8	U	U	U	E	VG	G-VG	VG	14 days
7	Fluopyram 10.9%											
11	Trifloxystrobin 13.1%											
7	Pydiflumetofen 7.0%	Miravis Neo 2.5 SE	13.7	13.7	U	U	U	E	VG-E	VG	G-VG	30 days
11	Azoxystrobin 9.3%											
3	Propiconazole 11.6%											
11	Pyraclostrobin 28.58%	Priaxor 4.17 SC	4.0-8.0	5	U	VG	U	VG	VG-E	VG	U ³	21 days
7	Fluxapyroxad 14.33%											
11	Pyraclostrobin 13.64%	Headline AMP 1.68 SC	10.0-14.4	10	U	E	E	E	VG	G	G-VG ³	20 days
3	Metconazole 5.14%											
11	Trifloxystrobin 32.3%	Stratego YLD 4.18 SC	2.0-5.0	4	VG	E	VG	E	VG	G	NL	14 days
3	Prothioconazole 10.8%											
3	Tetraconazole 7.48%	Affiance 1.5 SC	10.0-17.0	NA	U	G-VG	U	G-VG	G-VG	G	G ³	7 days
11	Azoxystrobin 9.35%											
3	Flutriafol 15.7%	Adastrio 4.0 SC	7.0-9.0	8	U	U	U	VG-E	U	VG	G-VG ³	30 days
11	Azoxystrobin 15.7%											
7	Fluindapyr 10.5%											

¹Fungicide application timing is extremely important and needs to be made near the onset of the tar spot symptoms. ²Harvest restrictions are listed for field corn harvested for grain. Restrictions may vary for other types of corn (sweet, seed, or popcorn, etc.), and corn for other uses such as forage or fodder. ³A 2ee label is available for several fungicides for control of tar spot, however efficacy data are limited. Check 2ee labels carefully, as not all products have 2ee labels in all states. This information is provided only as a guide. It is the applicator's legal responsibility to read and follow all current label directions. Reference in this publication to any specific commercial product is for general information only, and does not constitute an endorsement or recommendation by the CDWG. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer. Members or participants in the CDWG assume no liability resulting from the use of these products.



Scan this QR code to download the chart to the left.